

Pediatric CDAI

Abdominal pain

1



None	0
Mild	5
Moderate/Severe	10

Stool frequency (per day)

2



0-1 no blood	0
2-5 liquid or 1-2 semi-formed with blood	5
>5 liquid, gross blood/nocturnal diarrhea	10

General well-being


3



Very well	0
Below par	5
Very poor	10

Extraintestinal (5 points if 1, 10 points if >1)

4



Arthritis; Uveitis; E.nodosum; P. gangrenous;
Fever

5



Weight gain or stable	0
Involuntary stable or 1-9% weight loss	5
>9% weight loss	10

6



Height <1 channel decrease	0
Channel decrease 1-<2	5
>2 channel decrease	10

7



Abdominal mass	
None	0
Tenderness	5
Tenderness, guarding/definite mass	10



Pediatric CDAI


8



Perirectal disease

None	0
1-2 indolent fistulae, scant drainage	5
Active fistula, draining/abscess	10

9



Laboratory parameters

Use age-specific reference

Hematocrit %

- Normal
- Mild decrease
- Moderate/severe

0
2.5
5

Erythrocyte sedimentation rate

- <20
- 20-50
- >50

0
2.5
5

Albumin (gr/L)

- ≥ 3.5
- 3.1-3.4
- ≤ 3

0
5
10

SCORE

<10

10-30

>30

Remission

Mild activity

Moderate-severe

