IBD THERAPIES PREGNANCY & LACTATION

	Pregnancy	Lactation
Mesalazine		
Sulfasalazine		
Azathioprine		
Mercaptopurine		
Methotrexate		Passes into the milk so recommended to avoid
Infliximab		
Adalimumab		
Golimumab		
Certolizumab		
Vedolizumab		
Ustekinumab		
Risankizumab		
Mirikizumab		
Tofacitinib		Passes into the milk & manufacturers recommend to avoid it & at least 18 hours after the last dose
Filgotinib		Passes into the milk & exposure to newborn cannot be excluded
Upadacitinib		Very little data but likely to apply same concept than to other JAKs and avoidance is recommended
Ozanimod	Scarce data to be avoided	Very little data but the metabolites can be detected in the milk the manufacturers recommend to avoid
Etrasimod	Scarce data to be avoided	Not enough information is available on the use of etrasimod during breastfeeding

ECCO 2022 & AGA 2019 guidelines

Julsgard M et al. Lancet. 2023; 8 (8): 695-697 https://www.ema.europa.eu/en/documents/product-information/zeposia-epar-product-information_en.pdf https://ec.europa.eu/health/documents/community-register/2019/20191216146604/anx_146604_en.pdf



Low risk

To be avoided

No data-recommended to avoid

