

Subcutaneous Biologics cold chain cheat sheet

- Infliximab: stable 28 days $\leq 25^{\circ}\text{C}$.
- Adalimumab: stable 14 days $\leq 25^{\circ}\text{C}$.
- Certolizumab: stable 7 days $\leq 25^{\circ}\text{C}$.
- Golimumab: stable 30 days $\leq 25^{\circ}\text{C}$.
- Vedolizumab: stable 7 days $\leq 25^{\circ}\text{C}$.
- Ustekinumab: up to 30 days $\leq 30^{\circ}\text{C}$.
- Mirikizumab: stable 14 days $\leq 30^{\circ}\text{C}$.
- Risankizumab: stable 24 hours $\leq 25^{\circ}\text{C}$.
- Guselkumab: stable 30 days $\leq 25^{\circ}\text{C}$.

These drugs should generally be stored between 2°C - 8°C if the drug has been frozen $< 2^{\circ}\text{C}$ it normally needs to be discarded. If you have any doubts please discuss it with your pharmacist

This table is intended for informational purposes only and does not replace medical advice. Storage conditions may vary depending on specific formulations and local regulations. Always consult the product's official labeling and discuss any concerns with the prescribing physician or pharmacist. Individual clinical judgment and common sense should prevail in every case.

